

Analysis of NOAA and NASA Surface Water and Energy Balance Data for Water Resources Applications


David Toll (NASA/GSFC)*, Jiarui Dong (UMBC/GEST), Paul Houser (GMU/CREW), Kristi Arsenault (UMBC/GEST), Ana Pineiro (UMBC/GEST), Jeff Basara (U. OK) and Justin Monroe (U. OK)

*NASA/GSFC Hydrological Sciences Branch/614.3, Greenbelt, MD 20771 USA (David.L.Toll@nasa.gov, 301-614-5801)

Abstract

The primary objective of the study is to evaluate surface water and energy balance for water resources applications through study of NOAA North American Regional Reanalysis (NARR) data and North American Land Information System (NLDAS) Mosaic and Noah land surface model heat flux data. For reference we used *in situ* data from the Coordinated Enhanced Observing Periods (CEOP 3&4), the Oklahoma Mesonet sites, and the NASA-Reclamation ET tower sites in the Middle Rio Grande, New Mexico. In addition, the high spatial resolution (to 1km and finer) NASA Land Information System (LIS) is used to further study factors affecting modeled surface water and energy budget data for future improvements to NARR and NLDAS type systems. Analysis of results for the CEOP study sites show significant biases are found for the partitioning between sensible heat and latent heat fluxes during spring and summer seasons. We used LIS to assess possible factors and evaluated improvements using revised model forcing, model physics, and input parameters. Current work in the Middle Rio Grande is developing an Ensemble Kalman Filter (EnKF) capability using MODIS Land Surface Temperature with the Community Land Model (CLM-2) for improved heat fluxes. Oklahoma Mesonet data are used to evaluate NOAA NARR and NLDAS Mosaic and Noah data. Work to date has shown the NARR data tracks well the extreme wet to dry periods during the summer of 2003.

STUDY SITES



NLDAS & NARR DATA SET SUMMARY

- 1) NARR: North American Regional Reanalysis (NOAA/NCEP)
 - Coupled land-atmosphere model, 30 Pressure Levels, 196 Variables
 - Temporal coverage: 1979-2003
 - Spatial resolution is at 32 km
 - Output at every 3-hour frequency
- 2) NLDAS NOAA NCEP NOAH LSM & NASA/GSFC MOSAIC LSM
 - Real time and retrospective hourly, uncoupled land surface models
 - Spatially 1/8th-degree grid cell (~13km)
 - Every 1-hour frequency output from 10/1996 to present
 - UMD Pinker's corrected GOES-based forcing

NASA LAND INFORMATION SYSTEM






Figure Land Information System (LIS) is a high performance land surface modeling and data assimilation system capable of modeling global land-atmosphere interactions at spatial resolutions to 1km and finer. LIS was used to further investigate factors affecting surface water and energy budget data.

Assessment of the Model Energy Budget

Golden Day Sensible (SH) and Latent Heat Flux – All sites



Although Golden Days improved net radiation estimates (left); however, in the figure above large RMS errors are still evident in the estimates of the partitioning of sensible heat and latent heat fluxes. This is further investigated using the NASA Land Information System.


(CEOP Cont'd) Using 'LIS' to Evaluate Surface Heat Fluxes

We used the LIS runs at a 1KM spatial resolution with three models (Noah, Mosaic & NCAR CLM2). Experiments were designed through LIS configurations with the original NLDAS forcing and *in-situ* data. We checked some possible reasons causing the partitioning problems between sensible heat and latent heat fluxes from (1) the inaccurate estimates of model forcing data, (2) a model scheme and parameterization, and (3) the land cover specifications.


FORCING DATA



SURFACE ALBEDO



LAND COVER



Replaced NLDAS Forcing w/ In Situ Forcing

- Shortwave, Precipitation, Longwave (↓), Ta, Qa & P
- Most significant change from Shortwave Radiation
- Improved surface heat fluxes for all 3-models (SH+LH shown)
- Large SW Errors, April-August 2003 (RMS of 220 vs. 50 W/m²). NASA/GSFC Mosaic different from NCEP Noah.


Replaced "Observed" albedo (in situ) in LIS Mosaic model

- In situ albedo directly assimilated (inserted) for each step
- Snow albedo not in Mosaic but in Observed (see Spikes)
- Improved accuracy of heat fluxes (shown SH+LH+G)
- Current work inserting in situ soil moisture

Updated NLDAS Land Cover


- 3 (E4, E7 & E20) of 12 Sites did not match in-situ cover
- Updated Dominant UMD Land Cover in NLDAS
- Significant improvements in Heat Fluxes
- Mosaic and CLM2 are more sensitive to Land Cover

II. Middle Rio Grande LIS ET & Ensemble Kalman Filter MODIS ET Evaluation




Temperature

- Assimilate MODIS LST from Rapid Response System
- Available within 4 hours of overpass



Ensemble Kalman Filter (EnKF)

- Implement EnKF in CLM-2
- 1 Km Spatial Resolution
- Developing LIS Toolbox




LIS ET Summary

- April 2004 ET (mm)
- Community Land Model (CLM-2)
- Middle Rio Grande
- 1 km Spatial Resolution
- Will test ET Data Assimilation using Eddy Covariance test sites


III. Oklahoma Mesonet (OASIS Super Sites)

NARR Surface Heat Flux Comparison



Comparison of NARR Surface Heat fluxes to Norman Oklahoma OASIS Super Site shows a large contrast in latent heat flux versus sensible heat flux due to a wet June and a dry July. Note, days (June, n=11 and July, n=20) were selected to show stark contrast. The NARR was able to pick up much of the trends.

Noah LSM Heat Fluxes



Previous study showing comparison of Noah LSM with Mesonet data showing a Strong correspondence. "Golden Days". (Nemunaitis and Basara, 2004)

Summary

This study evaluates NOAA and NASA land surface water and energy balance data with the goal to assess factors causing uncertainty with the goal to provide improved and useful surface energy and water flux data for water resources applications.

Acknowledgements

This work was funded under the NOAA-CPA. Additional support was provided by the NASA-HQ Terrestrial Hydrology Program and the NASA-GSFC Earth-Sun Exploration Division.