

# Ice supersaturation and cirrus clouds in HIPPO Global Campaign #1-5

Minghui Diao, Mark Zondlo  
Civil and Environmental Engineering  
Princeton University



2DC, 260X 2DC ice probes, Dave Rogers, Al Cooper

RAF Technical and Ground Crews

HIPPO Global Campaign Science Team

NASA Earth System Science Fellowship;  
Funding of NSF – 0840732;  
MIRTHE ERC  
2012-March-12

Photo by Minghui Diao



# Outline

- **Motivation**
  - cirrus clouds climate effects in NH vs SH
  - Ice supersaturation in cirrus cloud formation
- **Instrument and dataset**
  - HIPPO 1-5
  - Water vapor; Temperature; Ice crystal number density
  - Uncertainties
- **Ice supersaturation (ISS) in HIPPO Global campaigns**
  - ISS probability distribution function (PDF) in NH vs SH
  - Pole-to-Pole latitudinal and vertical distribution of ISS
- **ISS and cirrus clouds formation**
  - Separate difference phases of cirrus clouds in HIPPO
  - Cirrus cloud formation mechanism in NH vs. SH
- **Conclusion**

Photo by Minghui Diao

# Motivation

- **Cirrus clouds (235-185K, up to 40% coverage)**
  - Climate effect
    - Large & uncertain effect (IPCC, AR4, 2007)  
Warming or cooling (Chen et al. 2000)  
Difference NH vs. SH, anthropogenic activities, inhibit or invoke?  
Microphysical properties (ice crystal number and size distribution)
- **Ice supersaturation (ISS)**
  - Birthplaces of cirrus clouds: **relative humidity with respect to ice (RHi) > 100%**
  - Anthropogenic aerosols indirect effect (CCN, IN, lower ISS; organic aerosol, higher ISS)
- **Challenges in observations**
  - **Remote sensing** >> microphysical scale
  - **Small scale observations** limited by spatial temporal coverage
    - NH vs. SH, lack of sampling  
INCA Campaign Prestwick 55N and Punta Arenas 55S (Ovarlez et al., 2000)
- **Unsolved questions:**
  - What is the global distribution of ISS by *in situ* observations in HIPPO?
  - Is there any difference in cirrus cloud formation mechanism between NH and SH?

Photo by Minghui Diao

# Instrumentations and dataset

- **Instruments**
  - **Water vapor**: the VCSEL hygrometer (accuracy 6%)
  - **Temperature**: Rosemount temperature probe ( $\pm 0.5$  K)
  - **Ice crystal number density**: 2DC and 260X 2DC probes (25  $\mu\text{m}$ , 10  $\mu\text{m}$ )
- **HIPPO 1 to 5 deployments**
  - HIPPO1 did not have ice measurements
- **Uncertainties**

Relative humidity with respect to ice ( $T \leq -40$  C)

$$RHi = e/e_s$$

- $e$ : water vapor partial pressure
- $e_s$ : saturated ice vapor pressure

Example of RHi uncertainty

Water vapor mixing ratio: 6%  
Temperature: 0.5 K





RHi 8-10%  
@ 233-205K



# Clear sky RHi distribution in NH and SH


Ovarlez *et al.*, 2000

1 Hz



SH higher frequency of ISS

Probability density function of RHi





NH RHi distribution shifts to higher ISS  
NH: 71 hrs, SH: 26 hrs


# Cloudy sky

## RHi distribution in NH and SH

Ovarlez *et al.*, 2000



Peaks at ~95% (NH) and ~94% (SH)




No obvious difference  
NH: 4 hrs, SH: 1hrs

SH higher frequency of ISS

# Clear sky


## Ice supersaturation (ISS) distribution in NH and SH

ISS magnitude



**ISS magnitude** between two hemispheres  
NH has higher ISS

ISS vertical distribution




**# of ISS** between two hemispheres  
NH has more # ISS in observations


# Clear sky

## ISS frequency density in NH and SH

Bin by 25 mb\*10 degrees



**Total RHi observations  
In clear sky**




**ISS frequency density** between two hemispheres  
NH has higher frequency of ISS for clear sky

# Cloudy sky

## ISS distribution in NH and SH


ISS magnitude



HIPPO1 has no ice probe

**ISS magnitude** between two hemispheres  
NH has higher ISS

ISS vertical distribution




**# of ISS** between two hemispheres  
NH has more # ISS in observations


# Cloudy sky

## ISS frequency density in NH and SH

Bin by 25mb\*10degrees




**Total RHi observations  
In cloud**



No obvious difference for in-cloud ISS frequency  
Limited cloud data in SH

# Ice supersaturated regions (ISSRs) and ice clouds

**ISSRs:** spatially continuous region where  $RHi > 100\%$ ,  
with or without ice crystals







ISS magnitude is highly related to the formation and evolution of cirrus clouds




**During which phase of cirrus cloud evolution does NH has higher ISS than SH? – Nucleation? Removal?**

**How can we separate out different evolution phases of cirrus clouds?**



# Spatial ratio between ISSRs and cirrus clouds

**Spatial ratio = A / B**






# Phase 1 Clear sky ISSRs



Mean RH<sub>i</sub> value of each bin





Total ISSR + Cirrus clouds  
NH 3966; SH 4147

NH has broader scope of RH<sub>i</sub>  
inside totally clear sky ISSRs


-> PDF of RH<sub>i</sub> in clear sky

# Phase 2+3+4 Cirrus cloud growth





Color Phase 1,2,3

Mean RHi value of each bin




Red: NH; Purple: SH  
Not much difference  
NH has boarder RHi  
scope during cirrus  
cloud growth.

# Phase 5 Cloud sedimentation and evaporation



Mean RHi value of each bin



Red: NH; Purple: SH  
NH has many low RHi in-cloud.

# Conclusions

- 1. Ice supersaturation in NH and SH with global *in situ* HIPPO data
  - PDF of RHi NH shifts to higher ISS than SH for both in-cloud and clear sky
  - ISS frequency density
    - Clear sky NH > SH
    - In-cloud no difference
- 2. Evolutions from ISSR to cirrus clouds
  - Proposed a scheme to separate evolution of cirrus clouds by *in situ*, quasi-Eulerian sampling
  - Link large scale RHi difference to cloud microphysics
- 3. Mechanism of cirrus cloud formation in NH and SH
  - Separate out new-born clouds from aged clouds
  - NH vs. SH have similar mean RHi value at each cloud evolution phase
  - NH has broader range of RHi at each phase

Future work:

Meteorology, local sampling bias, instrument uncertainties  
(START08 SID\_Num\_2H).

Large scale dynamics? Aerosol background? Pollution?

# Acknowledgement

## RAF Technical and Ground Crews HIPPO Science Team



Thanks!  
Questions?



## Fellowship and award

2008-2012	Princeton Francis Upton Fellowship
2009-2012	NASA Earth and Space Science Fellowship
2011	Princeton Environment and Climate Scholars Travel Grant
2010	Outstanding student paper award for AGU Fall Meeting, San Francisco
2009	Outstanding student paper award for AGU Spring Assembly, Toronto
2009	Travel Award to attend Water Vapor and the Climate System (WAVACS) summer school, France